

St. Luke's University Health Network Health History Requirements for Contingent workers

Vaccine	Vaccination	Serologic Immunity
MMR (required) **	Proof of 2 MMR vaccines, a minimum of 4 weeks apart	Positive IgG titer for Measles (Rubeola), Mumps, Rubella
Varicella (chicken pox) **	Proof of 2 doses of Varivax, a minimum of 4 weeks apart	Positive IgG titer for Varicella
Tdap (Adacel, Boostrix) **	Proof of one dose of Tdap (tetanus, diphtheria, pertussis) within the last 10 years	N/A
Influenza**	Proof of the annual seasonal flu vaccine Oct 1 – March 31(or longer based on influenza activity as determined by SLUHN)	N/A
Hepatitis B (<i>optional, strongly recommended</i>)	Three-dose series of HepB vaccines are recommended for healthcare workers that may involve exposure to blood or body fluids. (dose #1 now, dose #2 in one month, dose #3 approx 5 months after dose #2)	Immunizations encouraged if risk of exposure to blood & body fluids: declination must be signed if proof of immunity not included

** These vaccines are required unless proof of positive titer is available. Live virus vaccines (MMR & Varicella) affect the accuracy of tuberculin skin tests (TST). Please complete all TST tests (including having them read) prior to receiving the MMR or Varicella vaccines. (If TST is performed and read within 4-6 weeks after receiving a live vaccine, there is a risk for a false-negative result due to temporary immunosuppression caused by the live virus vaccine).

The annual seasonal flu shot is required. Individuals requesting a medical exemption will be provided a SLUHN medical exemption request form for completion by their medical provider. SLUHN will conduct a formal review to determine if the request will be approved.

Tuberculosis Testing Options to be completed within 90 days of starting at St. Luke's:

- **BAMT: (Blood Assay Mycobacterium Tuberculosis) - Quantiferon Gold blood test (QFT) or TSPOT blood test:** negative result and completion of a TB Questionnaire and annual TB Questionnaire thereafter.
- OR
- **2 step Tuberculin Skin Testing: *2 step test definition:** 2 step testing is 2 tests administered 1-3 weeks apart with results and requires 4 visits.
 - Step 1 includes: Test 1 placement date; Test 1 read date (48-72 hours to have site assessed),
 - Step 2 includes: **1-3 weeks later after Test 1 is read:** Test 2 placement date, Test 2 read date (48-72 hours to have site assessed)
- OR
- **2 TST dates (if done annually),** with most recent test within last 3 months and a prior test result from one year ago. Annual test date results = consecutive years.
- Notes:*
- **BCG Vaccinated:** Those who have received the BCG vaccine in the past must have BAMT or QFT completed and negative TB Questionnaire
 - **Past positive TST/PPD/IGRA:** Provide copy of positive TB test, Chest X-Ray results (must be done with the past year), documentation if treated. *BCG administration without TB testing is not acceptable.*
 - **New positive:** If TST/PPD +, order QFT. If QFT +, need negative Chest X-Ray results prior to start date.