

NETWORK EMPLOYEE HEALTH SERVICES(P) 484.526.4704 ♦ (F) 484.526.4201

(E) employeehealth@sluhn.org

TUBERCULOSIS EDUCATION - ONBOARDING

The following information is based on the TB Guidelines from the Centers for Disease Control and Prevention (CDC) for healthcare personnel (HCP) MMWR 5/17/19

ONBOARDING: New hires are screened for TB at the time of the pre-employment physical, including a symptom evaluation, TB blood test for those without documented prior TB disease or infection, and individual TB risk assessment.

ONGOING SCREENING: Annual TB education is recommended for all HCP including information on TB exposure risk.

POSITIVE TEST: Treatment is encouraged for all HCP with untreated latent TB infection (LTBI) unless medically contraindicated.

EXPOSURES: After a known exposure to potentially infectious TB disease, HCP should have a timely symptom evaluation, TB testing (if no prior positive result), and additional testing if indicated. Those with initial negative tests should be retested 8–10 weeks after exposure. HCP with documented prior TB infection or disease are not retested but should have further evaluation if a concern for TB disease exists.

RISKS: HCP might have risks for TB exposure that are not related to their work in the U.S., or they might have risks for TB progression after baseline testing that necessitate special consideration, when these risks are unrecognized, HCP might experience TB disease and then transmit TB to others. Therefore, health care facilities should educate all HCP annually about TB, including risk factors, signs, and symptoms; in addition to encouraging HCP to discuss any potential occupational or non-occupational TB exposure with their primary care provider, employee health, or occupational medicine clinician. Certain groups might be at increased occupational risk for TB exposure such as pulmonologists and respiratory therapists. Some work settings might have a higher risk such as emergency departments and bronchoscopy units. Factors influencing risk of exposure include the number of patients with infectious pulmonary TB who are examined in these areas and delays in initiating airborne isolation precautions.

TB FACTS: TB is a disease which usually affects the lungs, typically spread from person to person through the airborne route when the person with active pulmonary TB disease coughs, sneezes, or speaks. It is caused by a bacterium called Mycobacterium Tuberculosis (MTB).

SYMPTOMS: Include a feeling of weakness, weight loss, fever, night sweats, chest pain, and coughing up blood.

PPE: HCP caring for patients with TB need to use appropriate respiratory protection (N95 particulate respirator) which requires annual fit testing; including when caring for patients placed on airborne precautions and during a cough-inducing procedure like a bronchoscopy. Patients with suspected or confirmed TB need to be placed in private rooms with negative airflow or HEPA filter units, on Airborne Precautions, with the door closed...no one enters without wearing the appropriate mask.

Not everyone infected with TB becomes sick; there are 2 different TB related conditions; Latent TB infection (LTBI) and active TB disease...

LTBI: Latent TB infection is when the TB germs are in the body but the person is not sick or exhibiting TB symptoms because the germs are not active and can not spread to others; however, there is a risk they can develop active TB disease in the future – those with untreated LTBI should be encouraged to be treated.

TB DISEASE: Active pulmonary TB disease is when TB germs are actively multiplying and the person is sick with some or all symptoms of TB, they are capable of spreading the germs to others – those with active MTB disease need to be treated with medication, isolated until proven non-contagious, and take medication exactly as instructed to prevent becoming sick again or developing resistance.

RISK GROUPS: People who are most likely to get sick from TB are those with HIV, people who inject illegal drugs, babies and young children, elderly, people who were not treated correctly for MTB in the past, and people with chronic medical conditions.



Comments:

NETWORK EMPLOYEE HEALTH SERVICES(P) 484.526.4704 ♦ (F) 484.526.4201

(E) employeehealth@sluhn.org

TUBERCULOSIS SCREENING QUESTIONNAIRE - ONBOARDING

		•	
		Positive TB skin test (PPD or TST) Positive TB blood test (QFT or T-spot) Diagnosis of LTBI or TB disease Treatment for LTBI or TB disease Treatment for LTBI or TB disease Treatment for LTBI or TB disease NO	
•	Please o	wheek Ves or No to any NFW onset of the following symptoms	
•			
•			
•			
YES*	NO	History Review	
		Positive TB skin test (PPD or TST)	
		Positive TB blood test (QFT or T-spot)	
		Diagnosis of LTBI or TB disease	
YES*	NO	Symptom Description	
		Fatigue, malaise, extreme tiredness	
		Unexplained weight loss	
		Anorexia or loss of appetite	
		Fever of unknown etiology	
		Night Sweats unrelated to menopause	
		Productive cough lasting more than 2 weeks and <i>not from a cold</i>	
		Spitting or coughing up blood	
		Chest pain unrelated to cardiac history	
YES*	NO	Risk Assessment	
		Current or planned immunosuppression	
		Temporary or permanent residence (≥1 month) in a country with high TB rate	
*Addi	tional exp	planation:	
unders	stand tha	at if I have questions, develop any of these symptoms, or believe I was in contact with a	
Signature: Date:			
Emplo	yee Hea	lth: Complete this section if YES is checked above or QFT result is positive/indeterminate.	ant
Reviev	ved by: _	Date:	
Chest	X-ray:	Repeat QFT:	
QFT: _		Repeat QFT:	